

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

09/730,814

CAS ONLINE PRINTOUT

> d his

(FILE 'HOME' ENTERED AT 09:32:53 ON 27 SEP 2001)

FILE 'CAPLUS' ENTERED AT 09:33:08 ON 27 SEP 2001

E KYLE D/AU

E KYLE DON/AU

L1 55 S E4-E6

L2 0 S GOEHRING RIC/AU

E GOEHRING RIC/AU

L3 3 S E4-E5

L4 0 S L3 AND L2

L5 1 S L1 AND L3

SELECT RN L5 1

FILE 'REGISTRY' ENTERED AT 09:35:47 ON 27 SEP 2001

L6 24 S E1-E24

FILE 'REGISTRY' ENTERED AT 09:49:21 ON 27 SEP 2001

L7 STRUCTURE UPLOADED

L8 QUE L7

L9 19 S L8

L10 STRUCTURE UPLOADED

L11 3 S L10

L12 432 S L10 FUL

L13 STRUCTURE UPLOADED

L14 8 SEARCH L13 CSS SUB=L12 FULL


FILE 'CAPLUS' ENTERED AT 09:57:11 ON 27 SEP 2001

L15 5 S L14

=> d l13

L13 HAS NO ANSWERS

L13 STR



G1 Me, Et, n-Pr, i-Pr, n-Bu, i-Bu, s-Bu, t-Bu, H

G2 H, O


Structure attributes must be viewed using STN Express query preparation.

CAS ONLINE PRINTOUT

=> d bib abs hitstr 1-5

L15 ANSWER 1 OF 5 CAPLUS COPYRIGHT 2001 ACS
 AN 2001:416763 CAPLUS
 DN 135:33362
 TI Preparation of tertiary amino compounds having opioid receptor affinity
 IN Kyle, Donald; Goehring, R. Richard; Victory, Sam
 PA Euro-Celtique, S.A., Luxembourg
 SO PCT Int. Appl., 15 pp.
 CODEN: PIXXD2
 DT Patent
 LA English
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001039767	A1	20010607	WO 2000-US33047	20001206
	W:	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM			
	RW:	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR, BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG			
PRAI	US 1999-169396	P	19991206		
OS	MARPAT 135:33362				
GI					




AB The title compds. [I; R1 = a bond, alkenylene, etc.; R2, R3 = H, alkyl, alkenyl, etc.; R4 = a bond, alkenylene, etc.; R5 = H, 5-6 membered (hetero)aryl, cycloalkyl; R6 = alkyl, cycloalkyl, halo; n = 0-3], useful for the treatment of chronic and acute pain, were prepd. Thus, reacting benzaldehyde with 3-(dibutylamino)propylamine in the presence of NaBH₄, 3.ANG. mol. sieves in MeOH followed by amidation of the resulting I [R1 = (CH₂)₃; R2, R3 = Bu; R4 = a bond; R5, R6 = H] with phenylacetic acid in the presence of EDCI and DMAP in THF afforded I [R1 = (CH₂)₃; R2, R3 = Bu; R4 = COCH₂; R5 = Ph; R6 = H] which showed K_i of 40 nM against opioid receptor .mu. binding.


IT **343593-68-0P 343593-69-1P**
 RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); THU (Therapeutic use); BIOL (Biological study); PREP (Preparation); USES (Uses)
 (prepn. of tertiary amino compds. having opioid receptor affinity)

RN 343593-68-0 CAPLUS
 CN Benzenacetamide, N-[3-(dibutylamino)propyl]-N-(phenylmethyl)- (9CI) (CA INDEX NAME)

CAS ONLINE PRINTOUT



RN 343593-69-1 CAPLUS

CN Benzeneacetamide, N-[2-(dimethylamino)ethyl]-N-(phenylmethyl)- (9CI) (CA
INDEX NAME)

RE.CNT 1

RE

(1) Lowrie; US 3843657 A 1974 CAPLUS

L15 ANSWER 2 OF 5 CAPLUS COPYRIGHT 2001 ACS

AN 2001:115125 CAPLUS

DN 134:178566

TI Preparation of melanocortin-4 receptor binding compounds

IN Maguire, Martin P.; Dai, Mingshi; Vos, Tricia J.

PA Millennium Pharmaceuticals, Inc., USA

SO PCT Int. Appl., 215 pp.



CODEN: PIXXD2

DT Patent

LA English

FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	WO 2001010842	A2	20010215	WO 2000-US21327	20000804
	WO 2001010842	A3	20010816		
	W:				
	AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN,				
	CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR,				
	HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT,				
	LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU,				
	SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN,				
	YU, ZA, ZW, AM, AZ, BY, KG, KZ, MD, RU, TJ, TM				
	RW:				
	GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW, AT, BE, CH, CY,				
	DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, BF, BJ,				
	CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG				
PRAI	US 1999-147288	P	19990804		
	US 2000-223277	P	20000803		
OS	MARPAT 134:178566				
GI					




AB The title compds. of formula B-Z-E [wherein B = an anchor moiety; Z = a central moiety; E = an MC4-R interacting moiety], e.g. I [wherein P2, P3, and P4 = independently CH, CF, CCl, CBr, C(alkyl), C(alkoxy), C(CN), C(OH), or Cl; W1 = covalent bond or CH2; W2 = CH2, CHR3, or CR3R4; W3 = CH2, CHR5, or CR5R6; R = H or alkyl; Z1 = CH or covalently linked to Z2 to form a naphthyl ring; Z2 = CH, C(C.tplbond.CH), CCl, CBr, Cl, CF, or covalently linked to Z1 to form a naphthyl ring; Z5 = CH or C(OMe); R3-R6 = independently Me or Et], were prepd. and tested as melanocortin-4 receptor (MC4-R) binding agonists and antagonists. For example, .alpha.-tolunitrile in THF was added to a soln. of diisopropylamine in THF, which had been cooled to -78.degree.C and treated with BuLi. HMPA and 1-chloromethylnaphthalene in THF were added, the reaction cooled and stirred for 1 h, and the reaction quenched with H2O to give 2-(2-naphthalen-1-ylethyl)benzonitrile. Treatment with H2S and 1,3-diaminopropane, followed by heating to 80.degree.C for 72 h and work up, gave II. In a scintillation proximity assay (SPA) using high-throughput receptor binding screening, II showed exemplary inhibition of MC4-R. The invention compds., primarily 2-(2-arylalkylsulfanylphenyl)-4,5-dihydro-1H-imidazole and 1,4,5,6-tetrahydropyrimidine derivs., are useful in the treatment of disorders assocd. with wt. loss and pigmentation (no data).

IT 326486-16-2P

RL: BAC (Biological activity or effector, except adverse); SPN (Synthetic preparation); BIOL (Biological study); PREP (Preparation)
(inactive as MC4-R binding compd.; prepn. and high throughput MC4-R receptor binding screening of arylalkylsulfanylphenyl-substituted imidazoles and pyrimidines and analogs)

RN 326486-16-2 CAPLUS

CN 1,2-Ethanediamine, N,N-dimethyl-N'-(2-phenylethyl)-N'-(phenylmethyl)-(9CI) (CA INDEX NAME)



L15 ANSWER 3 OF 5 CAPLUS COPYRIGHT 2001 ACS

AN 1988:36842 CAPLUS

DN 108:36842

TI New synthetic "tricks". From aliphatic amines and amides to azides and/or how to convert RNHCOR' into RNHCOR" avoiding drastic hydrolyses

AU Garcia, Jordi; Vilarrasa, Jaume


CS Fac. Quim., Univ. Barcelona(III), Barcelona, 08028, Spain

SO Tetrahedron Lett. (1987), 28(3), 341-2

CODEN: TELEAY; ISSN: 0040-4039


CAS ONLINE PRINTOUT

DT Journal
 LA English
 OS CASREACT 108:36842
 AB Controlled redn. of N-alkyl-N-nitrosoamides to hydrazides followed by nitrosation and fragmentation affords azides in 80% overall yields, under mild conditions. This simple idea is the basis of methods for the conversion of alkylamines and N-alkylamides to alkyl azides, of RNHCOR1 into RNHCOR2 or RN(COR3)2, and of lactams into .omega.-azido esters or .omega.-azido acids.
 IT 112157-90-1P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (prepn. of)
 RN 112157-90-1 CAPLUS
 CN Benzeneacetic acid, 1-(phenylmethyl)hydrazide (9CI) (CA INDEX NAME)



L15 ANSWER 4 OF 5 CAPLUS COPYRIGHT 2001 ACS
 AN 1979:507817 CAPLUS
 DN 91:107817
 TI Phenylacetamide compounds
 IN Ajisawa, Yukiyooshi; Kamiyo, Tetsuhide; Saito, Takenao
 PA Kissei Pharmaceutical Co., Ltd., Japan
 SO Belg., 26 pp.
 CODEN: BEXXAL
 DT Patent
 LA French
 FAN.CNT 1

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	BE 870992	A1	19790201	BE 1978-190893	19781003
	JP 54055544	A2	19790502	JP 1977-119780	19771005
	JP 57005219	B4	19820129		
	GB 2006196	A	19790502	GB 1978-38905	19781002
	SE 7810421	A	19790406	SE 1978-10421	19781004
	DE 2843328	A1	19790419	DE 1978-2843328	19781004
	FR 2405240	A1	19790504	FR 1978-28383	19781004
	NL 7810053	A	19790409	NL 1978-10053	19781005
PRAI	JP 1977-119780		19771005		
GI					




I

AB Phenylacetic acids and acid chlorides reacted with ethylenediamines to give title compds. I [R = C1-3 alkoxy; R1, R2 (same or different) = C1-3 alkyl, or NR1R2 = heterocycllyl; R3 = H, C1-3 alkyl; R4 = H, C1-3 alkoxy; n, m (same or different) = 0, 1, or 2), which exhibited analgesic activity. Thus, a mixt. of 4-MeOC6H4CH2CO2H and PhCH2CH2NHCH2CH2NEt2 was heated 6-8 h at 130-40.degree. and worked up to give I [R = 4-MeO (n = 1),


CAS ONLINE PRINTOUT

R1 = R2 = Et, R3 = R4 = H, m = 2].
 IT 71108-98-0P
 RL: SPN (Synthetic preparation); PREP (Preparation)
 (prepn. of)
 RN 71108-98-0 CAPLUS
 CN Benzeneacetamide, N-[2-(diethylamino)ethyl]-N-(phenylmethyl)- (9CI) (CA
 INDEX NAME)



L15 ANSWER 5 OF 5 CAPLUS COPYRIGHT 2001 ACS
 AN 1971:99273 CAPLUS
 DN 74:99273
 TI Sigmatropic rearrangements of diazenes
 AU Baldwin, Jack E.; Brown, James Edward; Hoefle, Gerhard
 CS Dep. Chem., Massachusetts Inst. Technol., Cambridge, Mass., USA
 SO J. Amer. Chem. Soc. (1971), 93(3), 788-9
 CODEN: JACSAT
 DT Journal
 LA English
 AB Allylic hydrazines RN(NH2)CH2CH:CR1R2 are treated with HgO to give allylic
 azo compds. RN:NCR1R2CH:CH2 (I). I are obtained via diazenes
 RN+(:N-)CH2CH:CR1R2. Small amts. of hydrocarbons R2RCR1R2C:CH2, and
 CH2:CHCR1R2CR1R2CH:CH2 are also obtained but not from diazene
 intermediates.
 IT 31908-23-3
 RL: RCT (Reactant)
 (oxidn. of)
 RN 31908-23-3 CAPLUS
 CN Hydrazine, 1-benzyl-1-cinnamyl-, (E)- (8CI) (CA INDEX NAME)

Double bond geometry as shown.



=>